

## Central Valley Fall Run Chinook Salmon Spawner Escapement

### Escapement Number (#) 1990 - 1997

RIVER	1990	1991	1992	1993	1994	1995	1996	1997	Hist. Avg.*
	#	#	#	#	#	#	#	#	
Mokelumne	497	410	1,645	3,157	3,421	5,417	7,775	10,175	3,434
Stanislaus	403	485	560	1,068	1,380	671	217	1,500	5,376
Tuolumne	104	100	144	482	376	867	4,106	5,500	18,251
Merced	87	121	1,103	1,995	4,635	2,700	4,703	4,968	3,387
Cosumnes	no est.	0	0	0	0	no est.	no est.	no est.	921
American	11,565	25,273	10,927	37,442	36,219	74,518	75,033		37,875
Feather	31,125	38,160	40,542	40,694	51,435	71,776	64,575		46,310
Yuba	4,000	14,000	6,000	6,000	10,830	13,300	23,000		11,253
Battle Cr.	21,088	17,296	12,707	18,615	43,264	83,192	73,582		16,935
Sacramento	49,732	28,963	36,765	46,885	39,781	33,569	35,109		96,160
Sac. Tribs.	799	2,714	2,657	3,411	3,874	9,298	5,922		8,999

Data for San Joaquin tribe from Loudermilk (DFG); rest of Central Valley from DFG "Grandtab" database (Red Bluff).

Salmon escapement data is obtainable from several different sources. This data is the best obtainable; and may be different than other sources.

All data presented is provisional and subject of revision. Numbers do change after experts review more data sources.

Only years with actual numbers/estimates (including 0) were used to determine yearly averages.

\* = Historical average for San Joaquin tribe based on period from 1940-1997. For other Central Valley streams the period is 1940-1996.